


mi-eye 2™

CARE INSTRUCTIONS

A decorative graphic consisting of numerous blue dots of varying sizes, arranged in a curved, arc-like pattern that starts from the bottom left and curves towards the top right. The dots are scattered, with some larger and some smaller, creating a sense of movement and depth.

The Answer to a
Quicker Recovery

By choosing mi-eye 2™, you have taken the first step to getting back to your active life sooner than when choosing an MRI*. With the help of mi-eye 2™ you and your physician can now make a decision on what treatment is best for YOUR unique condition.

What's Next?

Following your mi-eye procedure, you will have to take care of the procedure site, as it is an open wound. You may also experience some aches, pains and drainage from the site. This is normal.

To make sure that it heals properly, follow the Care Instructions and any special instructions from you physician.

Follow Trice Medical on social media
and share your mi-eye 2™ story.



@tricemedical

Care Instructions

If you experience post-procedure aches or pains once the local anesthesia has worn off, you should first follow the **RICE** technique—**R**est, **I**ce, **C**ompression and **E**levation. If pain remains unresolved, you can take over-the-counter anti-inflammatory medications, such as an NSAID (ibuprofen, aspirin, etc.) or acetaminophen in accordance with labeled instructions. Be sure to follow your doctor's recommendations for the right course of at-home pain management, including which type of medication is best for you. If any chills, fever or warm sensation at the procedure site are experienced, or any other adverse effects, contact your physician as soon as possible.

Bandage

Prior to leaving the physician's office, a bandage was placed over the mi-eye procedure site. Any post-procedure bleeding should resolve quickly. If prescribed by your doctor, the wound can be left open to air within a day of procedure. If continued drainage from the wound is observed, you can re-apply a bandage to the site. Contact your physician if drainage appears abnormal or seems excessive.

Returning to Activity

Unless you experience any lingering symptoms, you can return to activity following a mi-eye procedure. Be sure to follow any recommendations for restricted activity your physician may have advised.

Don'ts

It is recommended to refrain from soaking the area until it has fully healed to prevent potential infection. For a period of time, as directed by your physician, you should refrain from the following:

- Bathtubs
- Hot tubs
- Swimming (pools, lakes, streams, oceans, etc)

mi-eye 2™ is for use in diagnostic and operative arthroscopic and endoscopic procedures to provide illumination and visualization of an interior cavity of the body through either a natural or surgical opening, and is available by prescription only. Talk to your doctor to learn more about mi-eye 2™

**Based on estimated timeframe for radiology testing and results, including follow-up consultation.*

